

Detailed analysis of η production in proton-proton collisions

A. Švarc, S. Ceci

*Rudjer Bošković Institute, Bijenička c. 54, 10000 Zagreb, Croatia,
E-mail: svarc@rudjer.irb.hr, ceci@rudjer.irb.hr*

Coupled-channel, multiresonance partial wave analysis (PWA), developed and used by Carnegie-Mellon-Berkeley analyses group (CMU-LBL 79)¹ has been recently very frequently used, and agreed as a potentially safe tool for extracting N^* resonance parameters^{2,3,4,5}. The newly formed subsection for resonance parameter analyses of Baryon Resonance Analysis Group (BRAG)⁶ has decided to re-evaluate relatively old analyses^{1,7}, what has been suggested even by their own authors recently⁸. BRAG has chosen three independent analyses which repeat the formalism suggested earlier, but with the use of new and improved data⁹. They have come to a certain level of agreement regarding the number of poles and their values. However, some of the "Cutkosky" like analyses have been dropped out^{10,11}, for technical reasons presumably.

However, one possible direction of analysis has been dropped altogether. The question arises whether the obtained PWA can be "relatively" safely taken as input to processes involving more than two particles in the final state. Temporarily forgetting the expected additional complications (initial and final state interactions, off-mass shell behavior of two body amplitudes, etc.) this work tends to estimate whether the present two body T-matrices can account for the observables of a three body process $pp \rightarrow pp\eta$, very carefully measured in Uppsala near the threshold¹². The special attention has been given to understanding the apparently inverted shape of the proton-eta differential cross section in the final state^{12,13}. The tendency of this work is not to improve the two body fit (which needs a lot of additional observables even to be semi-reliable,) but to see if the present T-matrices can explain the $2 \rightarrow 3$ body processes without drastic assumptions of the complications of the three body physics.

The first, and natural test of the reliability of the two body amplitudes appeared in the carefully measured total and differential cross section for the process $pp \rightarrow pp\eta$ ¹². We have developed a simple model based on the exchange of the lowest mesons depicted in the following figure:

Unfortunately, there is a number of models which claim to reproduce the results, but differ among themselves drastically^{14,16,15,17,18}, so it has been left to us to show that our model¹⁹, reduced to the assumptions of the mentioned models, gives a very similar result. The comparison is successfully made and will be shown elsewhere. The main idea of this presentation is to draw atten-




Figure 1: The meson-exchange mechanism of $pp \rightarrow pp\eta$ reaction. The initial and final state interactions are denoted as ISI and FSI respectively.

tion to the fact that the differential $p\eta$ cross sections in a three body process tends to show a different curvature when compared to the two body $\pi^-p \rightarrow \eta n$ process which should dominate the process¹². In spite of the additional uncertainties of the processes like ISI, FSI and off-mass shell extrapolation of two body amplitudes, the effects should be extremely high, and acting in the same direction in order to turn the slope of the differential cross section. The disturbing data are shown in Fig.2.

It is to be expected that the two $\rightarrow$ three body process is dominated by the two body proton-meson $\rightarrow$ proton η amplitude, in the vicinity of the threshold in particular. However, it turns out that even the shape of the differential cross sections of the impulse approximation two body process²⁰ and the measured 2 $\rightarrow$ 3 body processes are different. Let us just mention that only higher partial waves (like D_{13}) can account for the opposite curvature. Therefore, we are left with only two possibilities: either the ISI, FSI and off-mass shell effects of the higher order processes are responsible for the discrepancy, or the D_{13} partial wave is not confidently extracted in 2 $\rightarrow$ 2 body processes.

As it is obvious, the lowest partial wave in two body processes which can cause such a curvature are D_{13} , partial waves and they have been under close scrutiny at the Mainz workshop. Our feeling is that ISI, FSI and off-mass shell




Figure 2: The comparison of the experimental values (full dots with error bars) and the polynomial fit to them (dotted line) with $\pi^-p \rightarrow \eta n$ at the comparable energy (full triangles) with the Zagreb calculation (full thin line).

effects of higher order processes should be surprisingly strong to account for such a drastic change in the shape of the differential cross section. Therefore, there is an open possibility that something remains hidden in the D_{13} two body partial wave which we have not been able to detect in two body processes. The future goal is to investigate all suggested possibilities, and see whether the ρ exchanged meson domination in the hadron η production channel, which is quite an opened problem, can account for the apparent disagreement. The results of further research will be reported.

References

1. R.E. Cutkosky, C.P. Forsyth, R.E. Hendrick, and R.L. Kelly, *Phys. Rev. D* **20**, 2839 (1979).
2. D.M. Manley and E.M. Saleski, *Phys. Rev. D* **45**, 4002 (1992).
3. S. Capstick *et al*, *Phys. Rev. C* **59**, R3002 (1999).
4. C. Bennhold, e-Print Archive: nucl-th/0008023, to be published in the Proceedings of the 18th Indonesian National Physics Symposium.
5. T.P. Vrana, S.A. Dytman and T.-S.H. Lee, *Phys. Rept.* **328** 181 (2000).
6. BRAG on web, <http://cnr2.kent.edu/~manley/BRAG.html>
7. G. Höhler, in *Elastic and Charge Exchange Scattering of Elementary Particles*, edited by H. Schopper, Landolt-Börnstein, New Series, Group X, Vol.9, Part 2b, (Springer-Verlag, Berlin 1983).
8. G. Höhler, this conference
9. BRAG, this conference.
10. M. Batinić, I. Šlaus and A. Švarc, *Phys. Rev. C* **52**, 2188 (1995);
M. Batinić *et al*, *Physica Scripta* **58**, 15 (1998).
11. B. Krusche *et al.*, *Phys. Rev. Lett.* **74**, 3736 (1995);
B. Krusche *et al.*, *Phys. Lett. B* **397**, 171 (1997).
12. H. Calén *et al*, *Phys. Lett. B* **458**, 190 (1999).
13. B.M.K. Nefkens, this conference
14. J.M. Laget and F. Wellers, *Phys. Lett. B* **257**, 254 (1991).
15. G. Fäldt and C. Wilkin, TSL/ISV-96-0162 Uppsala University preprint.
16. T. Vetter *et al*, *Phys. Lett. B* **263**, 153 (1991).
17. E. Gedalin, A. Moalem and L. Razdolskaja, *Nucl. Phys. A* **634**, 368 (1998)
18. A.B. Santra and B.K. Jain, *Nucl. Phys. A* **634**, 309 (1998).
19. M. Batinić, A. Švarc and T.-S.H. Lee, *Physica Scripta* **56** 321 (1997).
20. S. Ceci and A. Švarc, work in progress.